of, 67, 151; elevation of, 67, 152; supposed protrusion of among secondary rocks, 69.

- Granitic mountains, aspect of, 59.
- veins shoot up into superincumbent rocks, 63; instance of at
- Mousehole in Cornwall, ib.; also at Glentilt, 64.
- Granular, composed of grains, 38.
- Gravel, beds of on the summits of detached hills, 354.

Green marl of the Paris basin, 229.

- sand, the arenaceous beds below chalk, so called, position of, 197; description of, 200; upper and lower green sand separated by a bed of stiff clay called galt, 201; the lower green sand generally ferruginous, ib.
- Greenstone, the diabase of the French, sometimes called trap, 99, 129.
- Grès rouge, grès des Vosges, grès bigarré, 163.
- Greywacke, or greywacke slate, German grauwacké, French traumate, 86; a coarse slate containing particles of other rocks and minerals, ib.; when the particles are very minute, passes into common state, ib.; when the fragments are numerous, and the slate scarcely perceived, resembles coarse sandstone or gritstone, ib.; described by the French as a transition sandstone, ib.; formation of, 86; conglomerate associated with, ib.
- Guadaloupe, skeleton of a woman found in the calcareous sandstone, 14.
- Gypseous marl and gypsum, detached hills on the banks of the Marne and Seine, 227; gypsum formation, ib.; organic remains in, ib.; bones of large quadrupeds found in, 228; bones of birds found in, ib.; freshwater shells in, separated from the marine shells by a bed of green marl, ib.
- Gypsum, or sulphate of lime, called also plaster stone and plaster of Paris, less abundant than carbonate of lime, 37; constituent parts of, ib.; accompanying rock salt in the Alps, anhydrous, 175, 176.

Hading, or dipping of a metallic vein, 286

- Hall, Sir James, his experiments to explain the formation of basalt, 146; on limestone and chalk, ib.; on sandstone, 210; his theory on the formation of valleys and on diluvian agency, 354.
- Halley, Dr., hypothesis, 3.
- Hastings sand, or iron sand, 192. Heber, Bishop, his account of the ele- Inclination of strata, 43.

phants of the Himmalaya Mountains,

- Height of mountains, table of. See Appendix, 376.
- Herbivorous quadrupeds, remains of in tertiary strata, 25.
- High Stile Mountain, 132; crater of. 133.
- Himmalaya or Himmaleh Mountains, ten thousand feet higher than Mont Blanc, 61; believed to be composed of secondary strata, 62; elephants of, with shaggy hair, 25.
- Hippopotamus, fossil remains of, common in England, France, &c., 331; tooth, cut of, 330.

Hone, or Whetstone slate, 85.

- Hornblende rock, called by the French amphibole, 36; melts easily into black glass, ib.; forms trap rocks, ib.; analyses of, ib.; granular and lamellar, 98.
- slate fibrous, and has a velvet lustre, 79; passes by gradation into serpentine, *ib.*; rare in England, but abundant in Scotland, 80; forms part of the principal mountain ranges in Europe, *ib*.
- Hornstone, or petrosilex, infusible without the addition of alkalies, 50.
- Hot springs. See Thermal waters.
- Human bones not discovered in a fossil state, 13, 25, 333; not more perishable than those of quadrupeds, note, 25.
- skeletons in caverns, intermixed with bones of extinct species of quadrupeds, 305, 307.
- Humboldt's account of the formation of new islands, 49; on volcanoes and the extent of volcanic fire, 252, 258, 260, 268.
- Hushing, used in Westmoreland for discovering beds of slate, 346.
- Huttonian theory respecting granite, 66; of metallic veins, 292; on the formation of valleys, 349.
- Hyena, bones of in Kirkdale cavern, 308.
- Hylaosaurus, or forest lizard, bones of discovered by Mr. Mantell, 197; unlike any known species, ib.

I. and J.

Jasper, 34, 79; beds of in the Apennines, 98.

Java, volcanic eruptions in, 265.

- Icthyosaurus, 24; skeleton, cut of, 181; conjectures respecting its being an inhabitant of the present ocean, 213.
- Jet, 119.
- Iguanodon, an enormous fossil herbiverous reptile, discovered by Mr. Mantell; teeth, cut, of, 195.

H.