


Fig. 45.



## FAMILY XII. CELLEPORIDÆ.

### 32. CELLEPORA,\* Otho Fabricius.

CHARACTER. *Polypidom calcareous, cellular, irregularly lobed or ramous, formed of urceolate cells heaped together or arranged in quincunx.—Polypes ascidian.*

1. C. PUMICOSA, *glomerous, rough, porous ; cells suborbicular, the mouth round with a produced marginal process.* J. Ellis.

PLATE XXXII. Fig. 1, 2, 3.

Porous Eschara, *Ellis*, Corall. 75, no. 7, pl. 27, fig. f, F, and 72, pl. 30, fig. d, D.—Tubipora verrucosa, *Lin.* Syst. edit. 10, 789.—Cellepora pumicosa, *Lin.* Syst. 1286. *Mull.* Zool. Dan. prod. 253. *Turt.* Gmel. iv. 640. *Berk.* Syn. i. 212. *Wern.* Mem. i. 560. *Turt.* Brit. Faun. 206. *Stew.* Elem. ii. 428, pl. 12, fig. 16, 17, copied from *Ellis*. *Lam.* Anim. s. Vert. ii. 170. 2de edit. ii. 256. Corall. 40. *Bosc*, Vers, iii. 147, pl. 3, fig. 3. *Flem.* Brit. Anim. 532. *Hogg's Stock*. 37. *Stark*, Elem. ii. 436. *Johnston* in *Trans. Newc. Soc.* ii. 267. *Templeton* in *Mag. Nat. Hist.* ix. 469.—C. verrucosa, *Lin.* Syst. 1286. *Fabric.* Faun. Grœnl. 434.—C. spinosa, *Turt.* Brit. Faun. 205.—Millepora pumicosa, *Pall.* Elench. 254. *Ellis* and *Soland.* Zooph. 135. *Turt.* Gmel. iv. 639. *Wern.* Mem. i. 560. *Stew.* Elem. ii. 428.—Flustra bullata? *Ellis* and *Soland.* Zooph. 16. *Stew.* Elem. ii. 436. *Turt.* Gmel. iv. 664—Le cellepore pouce et C. verruqueux, *Blainv.* Actinolog. 443-4.

Hab. Found on the stems of various corallines, on stones, and on the roots of Fuci, common.

The cells, by their conglomeration, form a porous friable calcareous

\* From *cella* a cell, and *περιττός*.