

CHAPTER II.

THE NATURAL FAMILIES OF THE CTENOPHORÆ.

SECTION 1. *Family characters in general among Ctenophoræ.*—As in all classes of the animal kingdom, the natural families of the Acalephæ, and those of the Ctenophoræ in particular, are characterized by particular patterns of form, determined by structural features bearing upon form. p. 187-190.

SECTION 2. *The natural families of the Ctenophoræ Eurystomæ.*—Three families may be distinguished in this sub-order: the Beroidæ proper, the Nescidæ, and the Rangiidæ. p. 190-193.

SECTION 3. *The natural families of the Ctenophoræ Saccatae.*—This sub-order embraces also three natural families: the Mertensidæ, the Cydippidæ, and the Callianiridæ. p. 193-198; while the Cestidæ, p. 198, constitute a sub-order by themselves. p. 292.

SECTION 4. *The natural families of the Ctenophoræ Lobatae.*—Five families may now be distinguished in this sub-order: the Euramphæidæ, the Bolinidæ, the Mnemiidæ, the Calyptumidæ, and the Oeyroidæ. p. 199-202.

CHAPTER III.

NORTH AMERICAN CTENOPHORÆ.

SECTION 1. *The genus Pleurobrachia and its species.*—Natural limits of the genus. Its form and structure; the bulk of the body consists of gigantic cells; their arrangement; the locomotive flappers; the circulation of fluids; the tentacular apparatus; the lasso-cells of the tentacles illustrated by Prof. H. J. Clark; other structural details. p. 203-248.

SECTION 2. *The genus Bolina and its species; with remarks on allied genera.*—Characters of the genus; comparison with Pleurobrachia; the ambulacral tubes and the locomotive flappers have not the same extent; the whole chymiferous system and its ramifications differ in different types of Ctenophoræ. p. 249-269. Genus Mnemopsis, p. 269.

SECTION 3. *The genus Idyia and other true Beroids.*—Difference between Beroe and Idyia as genera. The species belonging to each. Description of the species most common along our coast. Its structure, p. 270. Idyopsis, p. 288.

SECTION 4. *Tubular view of the Ctenophora known at present.*—Systematic enumeration of all the Ctenophoræ thus far described, with their principal synomyms, p. 289-296.

SECTION 5. *Geographical distribution of the Ctenophora.*—First attempt to define the faunæ of the sea by the Ctenophore found in different parts of the world; distinction between faunæ and zoological zones. p. 297-301.