

Eudendrium by Ehrenberg, but formerly united with Tubularia. It will, however, appear in the sequel, that there are marked differences, not only between these

- Tubularia ramosa*, Linn. ("Ellis, Corall., 31, t. 16, Fig. a, t. 17, Fig. a, A"), Fauna Suecica, Ed. altera, 1761, p. 539, No. 2229.
- " *trichoides*, Pallas, Elenchus, 1766, p. 84, No. 41.
- " *ramosa*, Linn., Systema Nat., 1767, 12th ed., p. 1302.
- " " Gmelin, Lin. Syst. Nat., 1788, p. 3831.
- " " Turton, Brit. Fauna, 1807, p. 210.
- " " Lamouroux, Bull. Soc. Phil., Paris, 1812, p. 184.
- " " Lamouroux, Polypiers Flexibles, 1816, p. 231.
- " " Lamarek, Animaux sans Vert., 1816, p. 110.
- " " Fleming, British Animals, 1828, p. 552.
- " " Bose, Hist. des Vers, 2de éd., 1830, III. p. 89.
- " " Blainville, Dict. Sc. Nat., 1830, Tom. 60, p. 435.
- " " Blainville, Manuel d'Actinologie, 1834-1836 (see p. 687 for the date), p. 470.
- " " Milne-Edwards, Lamk. An. sans Vert., 2de éd., 1836, II. p. 126.
- " *ramosa* and *ramosa*, Johnston, Brit. Zoöph., 1st ed., 1838, p. 116.
- " *ramosa*, Gould, Report on the Invertebrata of Massachusetts, 1841, p. 350.
- " *ramosa*, Dalyell, Rare and Remarkable Animals, 1847, Vol. I. p. 50, Pls. VI.-X.
- Fistularia ramosa*, Müller, Prodr. Zool. Danica, 1776, p. 254.
- ? *Fistularia ramosa*, Fabricius, Fauna Greenland., 1780, p. 441.
- Sertularia racemosa*, Cuvolini, Polipi Marini, Mem., 1785; and translated by Sprengel into German, 1813, Pl. VI. Figs. 1-7.
- " " Gmelin, Lin. Syst., 13th ed., 1788, p. 3854.
- " " Lamouroux, Hist. Polyp. Flexibles, 1816, p. 195.
- " *ramosa*, Blainville, Dict. Sc. Nat., 1830, Tom. 60, p. 445.
- " " Blainville, Manuel d'Actinologie, 1834-1836 (see p. 687), p. 480.
- Campanularia* (*C. racemosa* = *Sertularia racemosa*, Cuvolini), M.-Edwards, in Lamk. An. sans Vert., 2de éd., 1836, p. 134.
- Calamella* (*C. trichoides* = *Tubularia trichoides*, Pallas = *T. ramosa*, Linn.), Oken, Lehrbuch, 1815, III. p. 55.
- ? *Thoa* (*T. Savignii* = *Tubularia ramosa*, Pallas, fide Lamx.), Lamouroux, Hist. Pol. Flex., 1816, p. 210, Pl. VI. Fig. a, B C. (*Thoa Halecina* = *Halecium*, Oken).
- " " Lamouroux, Exposition Method., 1821, p. 14, Pl. LXVII. Figs. 5 and 6.
- Corymbogonium* (*Eudendrium capillare*, Alder), Allman, Annals and Mag. Nat. Hist., 1861, VIII. p. 171.
- ? *Corythamnium* (*Eudendrium bacciferum*), Allman, An. Mag. Nat. Hist., 1859, Vol. IV. p. 52.
- HIPPOCRENE Mertens.**
- Hippocrene, Mertens, Brandt, Prod. Descrip. Anim., 1835, p. 29.
- " Mertens, in Brandt, Mém. Acad. St. Pet., 6th ser., II., 1838, p. 392.
- " Forbes, An. Mag. Nat. Hist., 1841, VII. p. 82, Pl. I. Fig. 2.