

*Thuiaria Flem.<sup>1</sup>*—*Sertularia Lin.* (p. p.).

*Th. Thuia Flem.*—*Sertularia Thuia Lin.*—*Europe* (Ellis).—Also  
*Th. articulata Flem.*

*Halecium Oken.<sup>2</sup>*—*Thoa Lamx.*—*Sertularia Lin.* (p. p.).

*H. halecinum Oken.<sup>3</sup>*—*Thoa halecina Lamx.*—*Sertularia halecina Lin.*—*Europe* (Ellis).

*Grammaria Stimp.* and *Cryptolaria Busk*, appear to be related to  
Halecium.

<sup>1</sup> *THUIARIA Fleming*, 1828.

- Thuiaria*, Fleming, British Animals, 1828, p. 545.  
" Johnston, Brit. Zoöph., 1838, p. 137.  
" Johnston, Brit. Zoöph., 1847, 2d ed.,  
p. 83.  
" Alder, Catalogue Zoöph. Northumb., &c.,  
1857, p. 27.  
" Hincks, An. Mag. Nat. Hist., 1861,  
VIII, p. 255.

- Biseraria*, Blainville, Diet. Se. Nat., 1830, LX,  
p. 446.  
" Blainville, Manuel d'Actinologie, 1834–  
1836, p. 482.

- Cellaria (C. thuia)*, Lamarek, An. sans Vert., 1816,  
p. 139.

- Sertularia*, Linnaeus, Systema Naturae, 1767, 12th ed.,  
p. 1308.

- " Fabricius, Fauna Gronland., 1780, p. 444.  
" Solander and Ellis, 1786, p. 41.

- Corallina resiculata*, *cavile angulato rigido*, *ramis*  
*dense stipitatis et bifurcati*, *terminantibus*  
*denticulis evanide appressis*.—Ellis,  
Corallines, 1755, p. 10, Pl. 5, fig. b, B.

<sup>2</sup> *HALECIUM Oken*, 1815. The short, simple,  
symmetrical calyces of the sterile hydra, which are  
too small to allow the animal to retreat in them,  
and the one-sided fertile calyces, seem to indicate  
a distinct family.

- Halecium*, Oken, Lehrbuch, 1815, III, p. 91.  
" Schweigger, Handbuch, 1820, p. 426.  
" Johnston, Brit. Zoöph., 1847, 2d ed.,  
p. 58.  
" Alder, Catal. Zoöph. Northumb., &c.,  
1857, p. 20.

*Halecium*, Alder, An. Mag. Nat. Hist., 1859,  
III, p. 354.

- " Hincks, An. Mag. Nat. Hist., 1861,  
VIII, p. 254.  
" Hincks, Proc. British Association, 1858,  
p. 128.

*Thoa*, Lamouroux, Hist. Polyp. Flexibles, 1816,  
p. 210.

- " Lamouroux, Exposition Méthodique, 1821,  
p. 14.  
" Johnston, British Zoöph., 1828, 1st ed., p. 119.  
" Blainville, Diet. Se. Nat., 1830, LX, p. 452.  
" Blainville, Manuel d'Actinologie, 1834–1836,  
p. 488.  
" Milne-Edwards, in Lamk. An. sans Vert.,  
1836, 2d ed., p. 147.

*Sertularia (halecina)*, Linnaeus, Fauna Suecia, 1761,  
p. 540.

- " " Linnaeus, Syst. Nat., 1767,  
12th ed., p. 1308.  
" " Solander and Ellis, Zoöphytes,  
1786, p. 46.

*Corallina erecta*, *tubulosa*, *pennata*, *halecis spinac*  
*facie*.—Ellis, Corallines, 1755,  
p. 17, Pl. 10.

<sup>3</sup> *Halecium Beanii Johnston*, and *H. muricatum*  
*Johnst.*, seem to me to be generically distinct from  
*H. halecium*, but I have not the means of ascertaining  
their true characters. If they constitute  
distinct genera, the characters now assigned to the  
genus will require modifications. Van Beneden has  
figured the animal of *H. halecium*, Bull. Acad.  
Belg., 1847, Vol. XIV, p. 462. The fertile hydra  
is double-headed and tentaculated.