

Fig. 136.



Rhinoceros.

Rhinoceros leptorhinus; fossil from freshwater beds of Grays, Essex (see p. 153); penultimate molar, lower jaw, left side; two-thirds of nat. size.

Fig. 137.



Hippopotamus.

Hippopotamus Pentlandi, H. v. Meyer; from cave near Palermo (see p. 160); molar tooth; two-thirds of nat. size.

Fig. 138.



Pig.

Sus scrofa, Lin. (common pig); from shell-marl, Forfarshire; posterior molar, lower jaw, nat. size.

Fig. 139.



Horse.

Equus caballus, Lin. (common horse); from the shell-marl, Forfarshire; second molar, lower jaw.

a. Grinding surface, two-thirds nat. size.
b. Side view of same, half nat. size.

Fig. 140.



Tapir.

Tapirus Americanus; recent; third molar, upper jaw; nat. size.

Fig. 141.




a. b. Deer.

Elk (*Cervus alces*, Lin.); recent; molar of upper jaw.

a. Grinding surface.
b. Side view; two-thirds of nat. size.

Fig. 142.



a. d. Ox.

Ox, common, from shell-marl, Forfarshire; true molar upper jaw; two-thirds nat. size.

c. Grinding surface.
d. Side view; fangs uppermost.