

- BUZAREINGUES, GIROU DE**, inheritance of tricks, i. 450.
- CABANIS**, pears grafted on the quince, ii. 246.
- CABBAGE**, i. 341-344; varieties of, i. 341; unity of character in flowers and seeds of, i. 342; cultivated by ancient Celts, *ibid.*; classification of varieties of, *ibid.*; ready crossing of, i. 343, ii. 68, 76, 110; origin of, i. 343; increased fertility of, when cultivated, ii. 91; growth of, in tropical countries, ii. 266.
- CABOOL**, vines of, i. 353.
- CABRAL**, on early cultivation in Brazil, i. 329.
- CACTUS**, growth of cochineal on, in India, ii. 265.
- CÆSAR**, *Bos primigenius* wild in Europe in the time of, i. 84; notice of fowls in Britain, i. 258; notice of the importation of horses by the Celts, ii. 187.
- CAFFRE** fowls, i. 241.
- CAFFRES**, different kinds of cattle possessed by the, i. 91.
- “**CÁGIAS**,” a breed of sheep, i. 99.
- Cairina moschata*, i. 191.
- CALCEOLARIAS**, i. 388; ii. 129; effects of seasonal conditions on, ii. 263; peloric flowers in, ii. 338.
- CALDWELL**, J., sporting of sugar-cane, i. 410.
- “**CALONGOS**,” a Columbian breed of cattle, i. 92.
- CALVER**, Mr., on a seedling peach producing both peaches and nectarines, i. 362.
- CALYX**, segments of the, converted into carpels, ii. 387.
- CAMEL**, its dislike to crossing water, i. 190.
- Camellia*, bud-variations in, i. 401; recognition of varieties cf, ii. 238; variety in, hardiness of, ii. 299.
- CAMERON**, D., on the cultivation of Alpine plants, ii. 147.
- CAMERONN**, Baron, value of English blood in race-horses, i. 456.
- Campanula medium*, ii. 185.
- CANARY-BIRD**, i. 311; conditions of inheritance in, i. 465; hybrids of, ii. 20; period of perfect plumage in, ii. 53; diminished fertility of, ii. 145; standard of perfection in, ii. 179; analogous variation in, ii. 341.
- CANCER**, heredity of, i. 452, ii. 56.
- CANFIELD**, Dr., on horses with curled hair, i. 58; on feral horses in North America, i. 64.
- CANINE** teeth, development of the, in mares, ii. 310.
- Canis alopec*, i. 30.
- Canis antarcticus*, i. 20.
- Canis argentatus*, ii. 134.
- Canis aureus*, i. 30.
- Canis cancrivorus*, domesticated and crossed in Guiana, i. 23.
- Canis cinereo-variegatus*, i. 30.
- Canis fulvus*, i. 30.
- Canis ingæ*, the naked Peruvian dog, i. 24.
- Canis latrans*, i. 27; resemblance of, to the Hare Indian dog, i. 22; one of the original stocks, i. 26.
- Canis lupaster*, i. 25.
- Canis lupus*, var. *occidentalis*, resemblance of, to North American dogs, i. 21; crossed with dogs, i. 23; one of the original stocks, i. 26.
- Canis mesomelas*, i. 26, 30.
- Canis primævus*, tamed by Mr. Hodgson, i. 27.
- Canis sabbar*, i. 25.
- Canis simensis*, possible original of greyhounds, i. 34.
- Canis thaleb*, i. 30.
- Canis variegatus*, i. 30.
- CANNING**, A. S. G., the japanned peacock, i. 305-307.
- CANTERBURY** Bell, doubled by selection, ii. 185.
- CAPE** of Good Hope, different kinds of cattle at the, i. 91; no useful plants derived from the, i. 328.
- CAPERCAILZIE**, breeding in captivity, ii. 139.
- Capra ægagrus* and *C. falconeri*, probable parents of domestic goat, i. 105.
- CAPSICUM**, i. 396.
- CARDAN**, on a variety of the walnut, i. 379; on grafted walnuts, ii. 247.
- CARDOON**, ii. 7.
- Carex rigida*, local sterility of the, ii. 154.
- CARLIER**, early selection of sheep, ii. 188.