

- Chimborazo (Mt.), 26, 274, 290, 296
 China, 51, 84, 145; Cambrian in, 482; Lower Silurian, 522; Upper Silurian, 564; Devonian, 628; Carboniferous, 682, 698, 696; Cretaceous, 833
 China Sea, 927
 Chinat Mts., 874
 Chipola epoch (group), 884, 891, 899*, 917
 Chipola sands (fossiliferous), 890, 891
 Chirolepis, 417, 620; Canadensis, 618*; Trallii, 417*
 Chiropters, 918
 Chirotherium, 772*, 778, 774; Reiteri, 692
 Chirox, 917
 Chiton, 424§; Canadensis, 514; carbonarius, 690
 Chlamydothorium, 1004
 Chlorine, 63
 Chlorite, 68§, 89
 — argillite, 89§
 — rocks, 79§, 88, 84, 86, 87, 319, 449
 — schist, 89§
 Chloritic marl, 865
 Chlorophyll, 136
 Cholesterol peculiaris, 646
 Cholodus, 692
 Chomatodus, 692
 Chondrites Colletti, 688
 Chondrodite, 63, 67§, 79, 319, 447, 449, 450, 531
 Chondrodite limestone, 79§, 449, 450, 531
 Chonetes, 546*, 550, 552, 562, 579, 611, 621, 622, 642, 674, 700; cornutus, 546*, 550; Dalmanianus, 703; deflectus, 592; Flemingi, 685; Hardensis, 625, 628; hemisphaericus, 590, 592; Illinoisensis, 642*, 647; latus, 427*, 567; lepidus, 612, 620; lineatus, 590; mesolobus, 675*, 690; mucronatus, 592, 602; Novascoticus, 562; ornatus, 642*, 646; planumbonus, 646; scitulus, 612, 620; setigerus, 595*, 601, 620; striatellus, 567, 568
 Chonophyllum Niagarensis, 547*, 550
 Choristoceras, 771; Haueri, 774
 Chouteau limestone, 637, 646
 Chrestotes Danae, 691; lapidea, 691
 Christiania, 309
 Christianite, 136
 Christmas Island, 151 (height)
 Chrome-spinel, 88
 Chrysalidina gradata, 432*, 860*
 Chrysoberyl, 449
 Chrysocolla, 335
 Chrysodomus Stonei, 917
 Chrysolite, 67§
 Chrysolithic gabbro, 272; hornblende, 532; pyroxenite, 532
 — rocks, 88-89
 Chrysomelids, 771
 Cicada, 419
 Cidarid, 59, 641, 760, 779, 834, 840; Blumenbachii, 778*, 791; clavigera, 866; cretosa, 866; florigemma, 790, 791; splendens, 854; Texana, 837; vesiculosa, 866
 Cimolestes incisus, 853*
 Cimoliosaurus, 845
 Cimolos Island, 296 (volcanoes)
 Cincinnati, Ohio, 533
 — anticline. See Cincinnati uplift
 — beds, 492, 504, 506, 511, 514, 515, 516; characteristic species, 516
 — epoch, 494, 559
 — group, 489
 — Island. See Cincinnati uplift
 — uplift, 387§, 490, 494, 522, 527, 532-533, 537, 539, 540, 633; influence of, in the Upper Silurian, 571
 Cinders, volcanic. See Volcanic
 Cinnabar, 335
 Cinnabar, Mt., 829
 Cinnamomum, 837, 896, 921; ellipticum, 839; Mississippiense, 895*, 896; Scheuchzeri, 839
 Cinnamon, 921
 Cinulia, 861; avellana, 861*; pulchella, 854
 Cirripeds, 420*, 421§, 513 (earliest), 579, 720
 Cladiscites tornatus, 771*
 Cladodus, 692, 702; Clarki, 619*, 620; Fyleri, 619*, 620; Kepleri, 620; marginatus, 702; sinuatus, 619*; spinosus, 644*, 647
 Cladonia, 75; rangiferina, 75
 Cladopora labiosa, 592
 Cladoxylon mirabile, 621
 Claiborne epoch (group), 884, 885, 889, 891, 916
 — (Lower), 884, 885, 888, 890, 896, 897*, 915, 916
 — (Upper), 896, 897*
 Clam, 423, 424§
 Claosaurus, 847, 856; annectens, 844*, 845*, 847
 Clarion coal, 652; sandstone, 652
 Clastic rocks, 75§
 Clathropora flabellata, 514
 Clathropteris, 740, 750; rectiuscula, 749*
 Clava Chipolana, 917
 Clavilithes humerosus, 916; Mississippensis, 916; pachyleurus, 916; Penrosei, 916
 Clay, 76§, 80§, 81 (kinds), 134
 Clay-ironstone, 70§, 82
 Clay marls, 815, 821, 854
 Clay shale, 638, 748, 892; slate, 80§, 84
 Clayey layers, plication of, 208, 209*
 — rocks, 12, 313
 Clayton beds, 888
 Clayton Peak, 360*, 361
 Clear Creek limestone, 543, 559
 Clear-Fork beds, 660
 Clear Lake. See Borax Lake
 Cleavage in rocks, 92, 112*, 113*, 370
 Cleodora, 425*
 Clepsydrops, 687
 Clepsysaurus Pennsylvanicus, 753, 754*
 Cleveland shale, 606, 619, 620
 Clidastes, 826; iguanavus, 848; propython, 848; velox, 848*
 Clidophorus Pallasi, 707
 Cliffs, wearing of, 220, 221*
 Climacograptus, 514, 520; bicornis, 510*; Emmonsii, 470*; typicallis, 516
 Clinactichnites Fosteri, 479*; Wilsoni, 479*; Youngi, 479*
 Climatal changes, causes of, 253-257
 Climatal development, 1026
 Climate, effects on the work of rivers, 189
 —, Cambrian, 484; Carbonic, 711-712; Champlain, 940; Cretaceous, 872-873, 877; Eocene, 929; Glacial, 940, 943, 944; Lower Silurian, 524; post-Mesozoic, 875, 877; Paleozoic, 727; post-Paleozoic, 736; Permian, 737; Quaternary, 940; Tertiary, 921, 939; Triassic and Jurassic, 791, 792-793; Upper Silurian, 574
 Clinch Mountain sandstone, 538
 Clinkstone, 85§
 Clinometer, 100*§; use of, for measuring distant slopes, 28
 Clinton beds or epoch, 356, 410, 535, 540, 544, 549-550, 552, 563, 570, 572, 577
 — and Medina, British equivalent of, 563
 Cliona sulphurea, 158
 Clitambonites, 500; Americanus, 515
 Clupea, 862
 Clymenia, 614 (first American), 620, 626; levigata, 627; Neapolitana, 614*; Sedgwicki, 626*; undulata, 627
 Clymenia limestone, 627
 Clypeus Hugi, 428*
 Coahuila Valley, 200
 Coal, 62§, 124, 136, 143, 154, 485, 727, 775 (jet); analyses, 661, 662, 663, 713; formation of, from vegetable debris, 712-714; impurities, 663, 664; kinds, 661; origin, 71, 155, 653, 654, 655; plant-remains in, 653-655, 658, 663, 664; structure, 709, 710; vegetable material of: kinds and composition, 712, 713
 — in Carboniferous, 493; Carboniferous and Subcarboniferous, 634, 636, 639, 648, 661-664, 674, 693, 694*, 695, 696; Permian, 660, 684, 685, 698; Triassic, 742, 744, 745, 748, 755, 769; Jurassic, 775, 776; Cretaceous, 818, 820, 822, 825, 826, 827, 828, 829, 831, 857, 865, 872; Tertiary, 887, 892, 920, 922, 927
 — areas of N. America, 635, 825
 — fields of Europe, 693, 694, 696
 Coal-beds, bowlders in, 664, 709; burning of, 84, 266, 313 (changed to coke); metamorphic changes in, 315, 453; thickness of, 651, 652, 653, 656, 657, 658
 Coal-measures, dirt-beds of, 653, 658; false, 639
 — section of, near Nesquehoning, Pa., 649*; at Trevorton Gap, Pa., 650*